

Calibration of nitridation reaction efficiencies from plasma wind tunnel data and beyond

Anabel del Val

Collaborators:



(VKI) M. Capriati, B. Helber, A. Turchi, O. Chazot, T. Magin

(Inria / École Polytechnique) P. Congedo, O. Le Maître

(UC Boulder) T. Minton

(UMN) T. Schwartzentruber

State-of-the-art nitridation efficiencies differ by several orders of magnitude



- Extension to **high temperature data**
- Models **do not account for carbon injection** in BL
- Points result from **complex numerical-experimental approaches**
- **Different** facilities and model **approximations** involved
- **Diverse** landscape in **treatment of the data**

Calibration of nitridation reaction efficiencies from plasma wind tunnel data and beyond

Development 1

High surface temperature
data & deterministic model

Development 2

Stochastic model: experimental
& parametric uncertainties

Development 3

Model-form uncertainties

Development 4

Merging of molecular beam
& Plasmatron data

Conclusions and future outlook

Development 1

High surface temperature data & deterministic model

Development 2


Stochastic model: experimental & parametric uncertainties

Development 3


Model-form uncertainties


Development 4

Merging of molecular beam & Plasmatron data




- Graphite samples exposed to a **subsonic plasma flow**
- Four conditions at different **surface temperatures** recorded [2,200-2,600K]
- **Recession rates** and **CN radiative signature** measured during steady state





Deterministic inverse problem



$$\arg \min_c J(\boldsymbol{p}) = \|\boldsymbol{y}_{\text{obs}} - \boldsymbol{f}(\boldsymbol{X}_{\text{obs}}, \boldsymbol{c}, \boldsymbol{p})\|^2 + \mathcal{R}(\boldsymbol{X}_{\text{obs}}, \boldsymbol{c}, \boldsymbol{p})$$

Development 1

High surface temperature data & deterministic model

Development 2


Stochastic model: experimental & parametric uncertainties

Development 3

Model-form uncertainties

Development 4

Merging of molecular beam & Plasmatron data



Deterministic inverse problem

Different surface temperatures

Everything else...


$$\arg \min_c J(p) = \|\mathbf{y}_{\text{obs}} - \mathbf{f}(\mathbf{X}_{\text{obs}}, \mathbf{c}, \mathbf{p})\|^2 + \mathcal{R}(\mathbf{X}_{\text{obs}}, \mathbf{c}, \mathbf{p})$$

Recession rate

Reacting flow model
w/ surface reaction

γ_N^{CN}

- What if \mathbf{y}_{obs} is noisy?
- What if we don't really know parameters \mathbf{p} ?
- What if \mathbf{f} is wrong?



Deterministic inverse problem

$$\arg \min_c J(\mathbf{p}) = \|\mathbf{y}_{\text{obs}} - \mathbf{f}(\mathbf{X}_{\text{obs}}, \mathbf{c}, \mathbf{p})\|^2 + \mathcal{R}(\mathbf{X}_{\text{obs}}, \mathbf{c}, \mathbf{p})$$


- What if \mathbf{y}_{obs} is noisy? → Aleatory uncertainty
- What if we don't really know parameters \mathbf{p} ? → Epistemic uncertainty
- What if \mathbf{f} is wrong? → Model-form uncertainty

Development 3

Stochastic inverse problem

$$\mathbf{y}_{\text{obs}} = \mathbf{f}(\mathbf{Q}, \mathbf{X}_{\text{obs}}) + \mathbf{E}$$

$$\mathbf{Q} = \{\mathbf{C}, \mathbf{P}\} \quad \mathbf{Q} \sim \mathcal{P}(\mathbf{q}), \mathbf{E} \sim \mathcal{P}(\mathbf{e})$$








Bayesian formalism

$$\mathcal{P}(Q | y_{\text{obs}}) = \frac{\mathcal{P}(y_{\text{obs}} | Q) \times \mathcal{P}(Q)}{\mathcal{P}(y_{\text{obs}})}$$


Posterior distribution Likelihood function Prior distribution

How to efficiently combine models, experiments and stochastic inference methods to extract accurate nitridation models with estimation of their uncertainty?





$$\gamma_N^{\text{CN}} = 0.28^{+0.22}_{-0.21} \exp\left(\frac{8337.8^{+1608.2}_{-3308.3}}{T_w}\right)$$



Development 1

High surface temperature data & deterministic model

Development 2



Stochastic model: experimental & parametric uncertainties

Development 3

Model-form uncertainties


Development 4

Merging of molecular beam & Plasmatron data



$$\arg \min_{\mathbf{c}} J(\mathbf{p}) = \|\mathbf{y}_{\text{obs}} - \mathbf{f}(\mathbf{X}_{\text{obs}}, \mathbf{c}, \mathbf{p})\|^2 + \mathcal{R}(\mathbf{X}_{\text{obs}}, \mathbf{c}, \mathbf{p})$$

- What if \mathbf{f} is wrong? → **Model-form uncertainty**







$$\frac{\mathcal{P}(\mathbf{y}_{\text{obs}} | \mathcal{M}_0)}{\sum_i \mathcal{P}(\mathbf{y}_{\text{obs}} | \mathcal{M}_i)}$$

$$\frac{\mathcal{P}(\mathbf{y}_{\text{obs}} | \mathcal{M}_1)}{\sum_i \mathcal{P}(\mathbf{y}_{\text{obs}} | \mathcal{M}_i)}$$


$$\frac{\mathcal{P}(\mathbf{y}_{\text{obs}} | \mathcal{M}_n)}{\sum_i \mathcal{P}(\mathbf{y}_{\text{obs}} | \mathcal{M}_i)}$$

Denominator of Bayes' rule







More informative experiments can be proposed based on these insights




Set of reactions [1]:




Models:
 1. Molecular beam experiments



2. Plasmatron experiments




(Future: Surrogate model from CFD stagnation line simulations)



[1] Prata, Schwartzentruber, Minton. *AIAA Journal* 60:2, 627-640, 2022

Capriati, del Val, Congedo, Minton, Schwartzentruber, Magin. (*ongoing...*)



Conclusions and future outlook

Take home messages

Uncertainty modeling/stochastic inverse problems

- Uncertainty is a necessary component that should be modeled in addition to the physical problem of interest
- Incorporating uncertainties in inverse problems requires the overhaul of deterministic inverse techniques
- The Bayesian formalism computes an objective measure of uncertainty, informed by experimental data

Development of Bayesian inference frameworks for the stochastic calibration of graphite nitridation

- A stochastic methodology has been developed to incorporate uncertainties affecting the calibration of nitridation
- Recession rates and CN densities used jointly to infer nitridation efficiencies
- Model-form uncertainties incorporated through a Bayesian Model Averaging method

Finite-rate nitridation model calibrated using molecular beam and Plasmatron data

- Detailed mechanism can get information from both experiments at low and high pressures
- Calibrated model seems to fit well the experimental data
- Still early in the work but hope to have results with a CFD model soon

THANK YOU

Questions?



von KARMAN INSTITUTE
FOR FLUID DYNAMICS

UTOPIÆ

